

2017 Annual HMO HEDIS® Report

	Care Provided to BCBSIL HMO Members	BCBSIL HMO HEDIS Rate	2017 Quality Compass National Average (excluding PPO)
Effecti	veness of Care		
Prever	ntion and Screening		
•	Adult BMI Assessment	94%	77%
•	Childhood Immunization: o Combination 3 Rate: 4 DTaP, 3 IPV, 1 MMR, 3 HiB, 3 Hep B, 1 VZV, 4 PCV	80%	76%
	 Combination 10 Rate: Combination 3 plus 1 Hep A, 2-3 RV, 2 Influenza 	50%	51%
•	Breast Cancer Screening	74%	73%
•	Cervical Cancer Screening	80%	74%
•	Colorectal Cancer Screening	71%	62%
•	Chlamydia Screening in Women - Age 16-24	43%	48%
•	Immunizations for Adolescents: o Combination 1 Rate: 1 Meningococcal, 1 Tdap/Td o Combination 2 Rate: Combination 1 plus HPV	83% 15%	77% 15%
•	Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents: o BMI Percentile o Counseling for Nutrition o Counseling for Physical Activity	86% 84% 77%	65% 61% 56%
Respir	atory Conditions		
•	Appropriate Testing for Children with Pharyngitis	83%	84%
•	Use of Spirometry Testing in the Assessment and Diagnosis of COPD	38%	42%
•	Pharmacotherapy Management of COPD Exacerbation o Systemic Corticosteroid o Bronchodilator	82% 85%	70% 77%
•	Medication Management For People With Asthma o Total - Medication Compliance 75% Asthma Medication Ratio - Total	49%	48%
Cardio	Astrima Medication Ratio - Lotal	82%	78%
		600/	620/
•	Controlling High Blood Pressure Total	69% 83%	62% 84%
Diabat	Persistence of Beta-Blocker Treatment After a Heart Attack	03%	04%
Diabet			
•	Comprehensive Diabetes Care o Hemoglobin A1c (HbA1c) Testing o HbA1c Control (<8.0%) o Eye Exam (Retinal) Performed o Medical Attention for Nephropathy	91% 64% 67% 90%	91% 56% 54% 90%
	 Blood Pressure Control < 140/90 mmHg 	73%	62%

Care Provided to BCBSIL HMO Members	BCBSIL HMO HEDIS Rate	2017 Quality Compass National Average (excluding PPO)
Effectiveness of Care		
Musculoskeletal Conditions		
Disease Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis	89%	88%
Medication Management		
Annual Monitoring for Patients on Persistent Medications – Total	78%	83%
Behavioral Health		
Antidepressant Medication Management		
 Effective Acute Phase Treatment 	64%	67%
 Effective Continuation Phase Treatment 	47%	51%
 Follow-Up Care For Children Prescribed ADHD Medications 		
 Initiation Phase 	39%	40%
Continuation & Maintenance Phase	45%	47%
Metabolic Monitoring for Children and Adolescents on Antipsychotics	28%	36%
Access/Availability of Care		
 Children and Adolescents' Access To Primary Care Practitioners 		
o 12-24 months	95%	98%
 25 months to 6 years 	87% 89%	92% 92%
7 to 11 years12 to 19 years	89%	92%
 12 to 19 years Initiation and Engagement of Alcohol and Other Drug Dependence 	0370	30 /0
Treatment		
Initiation of Treatment	35%	33%
 Engagement of Treatment 	10%	11%
Prenatal and Postpartum Care		
Timeliness of Prenatal Care	93%	85%
o Postpartum Care	82%	74%
Adults' Access To Preventive/Ambulatory Health Service – Total	93%	94%
Use of First-Line Psychosocial Care for Children and Adolescents on	59%	58%
Antipsychotics (Total)		
Utilization		
Well-Child Visits in the First 15 Months of Life - (6+ Visits)	71%	79%
Well-Child Visits In The Third, Fourth, Fifth and Sixth Years of Life	70%	77%
Adolescent Well-Care Visits	48%	48%
Overuse/Appropriateness		
Non-Recommended Cervical Cancer Screening in Adolescent Females	2%	2%
Appropriate Treatment for Children With URI	87%	88%
Avoidance of Antibiotic Treatment in Adults with Acute Bronchitis	23%	30%
Use of Imaging Studies for Low Back Pain	71%	74%
Measures Collected Through the CAHPS® Health Plan Survey	, .	
Flu Vaccinations for Adults Ages 18 – 64	49%	48%

Results are rounded to the nearest percentage. $\label{eq:hebbs} \textbf{HEDIS}^{@} \text{ is a registered trademark of the National Committee for Quality Assurance (NCQA).} \\ \textbf{CAHPS}^{@} \text{ is a registered trademark of the Agency for Healthcare Research and Quality (AHRQ).} \\$